



BEST AVAILABLE COPY

U.S. PATENT & TRADEMARK OFFICE
SEP 17 2005
I P E JC79



D





FIG. 1




FIG. 2





FIG. 3

Average # of tubes/well



HUVEC




FIG. 4A

IFN- γ assay





FIG. 4B


IP-10 assay

FIG. 4D

Average # of tubes/well



Average # of tubes/well



Average # of tubes/well




FIG. 5A

FIG. 5B

- PBS
- Endostatin (315 ng)
- ▨ Anti-IL-22R1 (315 ng) + Endostatin (315 ng)
- ▨ IP-10 (134 ng)
- ▨ IP-10 (134 ng) + Anti-IL-22R1 (134 ng)




FIG. 6A





FIG. 6C






FIG. 6D

**Number of CD31
Positive vessels/fied**





FIG. 6E

Study design

FIG. 7

Cohort #	# of patients	Viral dose (particles)	biopsy time (post inj.)
1	1	2×10^{10}	24 hrs
2	1	2×10^{11}	24 hrs
3	3	2×10^{12}	24 hrs
4	3	2×10^{12}	48 hrs
5	3	2×10^{12}	72/96 hrs
6	3	2×10^{12} (divided doses)	48 hrs
7	5	2×10^{12}	Core bx @ 30 days
8	5	2×10^{12} (2x/wk x 3 wks)	Core bx @ 30 days

FIG. 8



Q: Average pre-treat signal
is 4.4E+02, n=8

24 hrs 48 hrs 96 hrs

(Pre-treat MDA-7 IHC – all negative)

<u>Pt</u>	<u>Time (hr)</u>	<u>MDA-7 ctr</u>	<u>MDA-7 periph</u>	<u>TUNEL ctr</u>	<u>TUNEL periph</u>
1	24	20%	0 %	20 %	0 %
2	24	30 %	5 %	70 %	10 %
3	24	75 %	40 %	50 %	30 %
4	24	35 %	5 %	n.d.	n.d.
5	24	50 %	25 %	25 %	17 %
6	48	>60 %	20 %	40 %	17 %
7	48	60 %	5 %	70 %	10 %
8	48	20 %	0 %	5 %	0 %
9	96	50 %	25 %	80 %	25 %
10	96	90 %	0 %	35 %	0 %

FIG. 9

MDA-7 induced apoptosis < 50%

Kinetics of Serum Cytokine Response to INGN 241




FIG. 10

FIG. 15

Serum Cytokine Response to Intratumoral INGN 241 Treatment.

<u>Cohort</u> (dose)	<u>No. pts</u> <u>tested</u>	<u>No. positive (mean % peak increase)</u>				
		<u>IL-6</u>	<u>IL-10</u>	<u>IFNγ</u>	<u>TNFα</u>	<u>GM-CSF</u>
1 (2x10 ¹⁰)	1	1 (298%)	1 (291%)	1 (>1,000%)	0	1 (150%)
2 (2x10 ¹¹)	1	1 (143%)	1 (599%)	1 (281%)	1 (864%)	0
3 (2x10 ¹²)	3	1 <td>1 (134%)</td> <td>1 (317%)</td> <td>2 (345%)</td> <td>1 (100%)</td>	1 (134%)	1 (317%)	2 (345%)	1 (100%)
4 (2x10 ¹²)	3	3 <td>3 (221%)</td> <td>1 (173%)</td> <td>2 (71%)</td> <td>1 (120%)</td>	3 (221%)	1 (173%)	2 (71%)	1 (120%)
5 (2x10 ¹²)	3	3 (640%)	3 (400%)	1 (317%)	2 (156%)	0
7 (2x10 ¹²)	7	5 (604%)	6 (387%)	3 (860%)	3 (255%)	0
Total	18	14	15	8	10	3

FIG. 11

Level of Increased CD8+ T cell Frequency in Pts who Received Intratumoral INGN 241.

<u>Cohort (dose)</u>	<u>No. pts tested</u>	<u>No. with increased CD8+ T cells</u>	<u>Mean % CD3+CD8+ T cells¹</u>		
			<u>Pre- treatment</u>	<u>Post- treatment @</u>	<u>Day Post-treat.</u>
1 (2×10^{10})	1	0	29%	27%	
2 (2×10^{11})	1	1	32%	44%	Day 15
3 (2×10^{11})	3	3	33%	59%	Day 15
4 (2×10^{12})	3	1	35%	44%	Day 30
5 (2×10^{12})	3	1	22%	29%	Day 15
7 (2×10^{12})	7	2	30%	40%	Days 15-30
Total	18	8	31%	44%	Day 15

FIG. 12

Increase in Peripheral Blood CD8+ T Cells Following intratumoral INGN 241 Injection




FIG. 13




FIG. 14





FIG. 15



Days after Infection

FIG. 16





FIG. 17



Cell Cycle Phase

FIG. 18

Percent Survival (%)




FIG. 19B

Percent Survival (%)




FIG. 19A

FIG. 19D




FIG. 19C




FIG. 20B






FIG. 20A



TUNEL Positive (%)



TUNEL Positive (%)






FIG. 21



H1299



A549

FIG. 22







FIG. 23

Radiation Dose(Gy)




A549

Radiation Dose(Gy)



H1299

FIG. 24






FIG. 25




FIG. 26

FIG. 27




FIG. 28

	<u>IL-6</u>	<u>IFN-γ</u>	<u>IL-10</u>	<u>GM-CSF</u>	<u>S.Fas</u>	<u>IL-1b</u>	<u>TNF-α</u>	<u>pSTAT3</u>
Ad-	+	++	+					
<i>mda7</i>			decrease	+/-	++			
MDA-7	+	+++	decrease	n.d.	n.d.	n.d.	n.d.	++

+: > +++: increased cytokine secretion compared to Ad-luc control.

-: no change; n.d.: not done.

Cytokine levels were evaluated using ELISA from cultured cell supernatants taken 24-72 hr post treatment.

pSTAT3 results were analyzed 4hr after addition of MDA-7 protein and 24 hr after Ad-mda7 treatment using IHC.

Effect of Ad-md_A7 on A549 Lung Metastasis




FIG. 29




FIG. 30




FIG. 31

A




FIG. 32A

B



FIG. 32B




FIG. 32C






FIG. 33

FIG. 34





MDA-7/TUNEL correlation p < 0.01

FIG. 35




FIG. 36




FIG. 37




FIG. 38




FIG. 39

Av. copies/cell: 0.001

740

900

1

0.04

median # DNA copies / ug DNA




FIG. 40




FIG. 41

Single Dose (2E12 VP)

Patient Primary Histology # injections Clinical Activity (Y/N) Comment

Patient	Primary Histology	# injections	Local	Remote	Comment
71	Renal Cell Ca	1	N	N	
72	TCC Bladder	1	N	N	Withdrew prior to day 30
73	Melanoma	1	N	N	
74	Colo-Rectal	1	N	N	
75	SCCHN	1	N	N	
76	SCCHN	1	N	N	Withdrew prior to day 30
77	SCCHN	1	N	N	

Repeat Dose (2E12 VP Biweekly x3)

81	AdenoCa	3	-	Withdraw
83	Melanoma	12	Y	CR
84	Melanoma	12	N	Stable Disease
85	SCCHN	6	Y	Central necrosis
86	NSCLC	3	-	Not eval
87	SCCHN	6	Y	On study
88	Melanoma	6	Y	Pending

* Three lesions treated consecutively: PCR in 1st, cInCR in 2nd, regression in 3rd
** marked erythema around remote cutaneous lesion

FIG. 42




FIG. 43




FIG. 44





FIG. 45

48 hrs after infection





FIG. 46

H1299

I- β Ba →

Ractin →



IC 50 value
Ad-mda7

clone	($\times 10^3$ vp/cell)
neo	3.27 ± 1.1
clone1	2.58 ± 1.25
clone3	1.25 ± 0.3





FIG. 47






FIG. 49A




A

FIG. 49B




FIG. 49C




FIG. 49D





FIG. 50

Cell Viability MDAH 2774



Cell Viability (OVCA 420)





FIG. 51

Cell cycle analysis of MDAH 2774 at 72hrs



Cell Cycle analysis of OVCAR-420




FIG. 52

Cell Survival in MDA-MB-486





FIG. 53

A



B




FIG. 54




FIG. 55

Apoptotic Index

B






FIG. 56

A**B****C**

L-8



FIG. 57

Micro Vessel Density





FIG. 58

A


HUVECs

Survival Fraction (%)

**C**


HUVECs

Survival Fraction (%)

**B**

HUVECs

Survival Fraction (%)

**FIG. 59**

A





FIG. 60

B



Targeting Plasmid Constructs




FIG. 61

ER Targeting of MDA-7 Blocks Colony Formation




FIG. 62

R-targeted MDA-7 is pro-apoptotic




FIG. 63

Inhibition of human ovarian cancer cell proliferation by Ad-mda7




FIG. 64

Days after infection

Induction of cell cycle arrest by MDA-7 (MDAH 2774)

	G1	S	G2M	Apoptosis Sub G0
24 hrs				
PBS	30.55	34.55	34.9	2.05
Ad.Luc	32.95	36.6	30.45	1.41
Ad.mda-7	27.45	29.75	42.85	4
48 hrs				
PBS	40.8	32.6	26.6	0.955
Ad.Luc	28.75	32.45	38.8	0.84
Ad.mda-7	18.3	78	54.5	10.36

FIG. 65A

Ad-md α 7 induced G2/M cell cycle arrest in OVCA 420




FIG. 65B
OVCA-420

Induction of apoptosis due to overexpression of MDA-7 (MDAH2774)




FIG. 66

FIG. 67A




FIG. 67B




FIG. 68




FIG. 69




FIG. 70




FIG. 71A




FIG. 71B




FIG. 71C





FIG. 72






FIG. 74

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- BLACK BORDERS**
- IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- FADED TEXT OR DRAWING**
- BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- SKEWED/SLANTED IMAGES**
- COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- GRAY SCALE DOCUMENTS**
- LINES OR MARKS ON ORIGINAL DOCUMENT**
- REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.